

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

ORT1273

WHAT IS CLAIMED IS:

1. An isolated and purified nucleic acid molecule that encodes protease D-G protein, said nucleic acid molecule comprising a member selected from a group consisting of:
 - 5 (a) a nucleic acid molecule encoding a protein having at least a 70% identity to a polypeptide comprising amino acids 1 to 435 encoded by SEQ ID NO:2;
 - (b) a nucleic acid molecule encoding a protein having at least a 70% identity to a polypeptide comprising amino acids 1 to 292 encoded by SEQ ID NO:9
 - (b) a nucleic acid molecule which is complementary to either one of the
 - 10 polynucleotides (a) or (b) ;
 - (c) a nucleic acid molecule comprising at least 15 sequential bases of either one of the polynucleotides (a), (b), or (c); and
 - (d) a nucleic acid molecule that hybridizes under stringent conditions to either one of the polynucleotide molecules of (a) or (b).
- 15 2. The nucleic acid molecule of claim 1 wherein the polynucleotide is RNA.
3. The nucleic acid molecule of claim 1 wherein the polynucleotide is DNA.
- 20 4. The isolated and purified nucleic acid molecule of claim 1, having a nucleotide sequence selected from a group consisting of: (SEQ.ID.NO.:1), (SEQ.ID.NO.:8) and functional derivatives thereof.
5. The isolated and purified nucleic acid molecule of claim 1, wherein said nucleic
- 25 acid molecule is genomic DNA.
6. An expression vector for expression of a protease D-G protein in a recombinant host, wherein said vector contains a nucleic acid sequence encoding proteolytically active protease D-G protein and functional derivatives thereof.

30

ORT1273

7. The expression vector of claim 6, wherein the expression vector contains a nucleic acid molecule encoding protease D-G protein, having a nucleotide sequence selected from a group consisting of: (SEQ.ID.NO.:1); (SEQ.ID.NO.:8); and functional derivatives thereof.

5

8. The expression vector of claim 6, wherein the expression vector contains genomic DNA encoding protease D-G protein.

9. A recombinant host cell containing the expression vector of claim 6.

10

10. The recombinant host cell of claim 9, wherein said nucleic acid molecule has a nucleotide sequence selected from a group consisting of: (SEQ.ID.NO.:1); (SEQ.ID.NO.:8); and functional derivatives thereof.

15

11. The recombinant host cell of claim 9, wherein said cloned nucleic acid molecule is genomic DNA.

12. A protein, in substantially pure form having protease D-G proteolytic activity.

20

13. The protein according to claim 12, having an amino acid sequence selected from a group consisting of: (SEQ.ID.NO.:2), (SEQ.ID.NO.:9) and functional derivatives thereof.

25

14. A monospecific antibody immunologically reactive with protease D-G protein.

15. A process for expression of protease D-G protein in a recombinant host cell, comprising:

30 (a) transferring the expression vector of Claim 6 into suitable host cells; and

ORT1273

(b) culturing the host cells of step (a) under conditions which allow expression of the protease D-G protein from the expression vector.

16. A method to detect test compound modulation of proteolytic activity comprising
5 the steps;

a) incubating a test compound, proteolytically active protease D-G protein, and a labeled substrate for sufficient time to produce a detectable product as a result of proteolytic activity upon the labeled substrate; and

b) measuring a change in the quantity of product as a result of test compound
10 modulation of protease D-G proteolytic activity on the labeled substrate when compared to protease D-G proteolytic activity on the labeled substrate in the absence of test compound.

17. The method of claim 16 wherein the labeled substrates comprises a detectable
15 label selected from a group consisting of a radiolabeled atom, at least one fluorescent molecule, and a colorimetric molecule.

18. The method of claim 17 wherein the substrate is labeled with two fluorescent
20 molecules, and the detectable molecule is detected by fluorescent resonant energy transfer.

19. A compound active in the method of Claim 16, wherein said compound is a modulator of protease D-G proteolytic activity.

20. A compound active in the method of Claim 16, wherein said compound is
25 an agonist or antagonist of protease D-G proteolytic activity.

21. A compound active in the method of Claim 16, wherein said compound is a modulator of expression of protease D-G.

30

ORT1273

22. A pharmaceutical composition comprising a compound active in the method of Claim 16, wherein said compound is a modulator of protease D-G proteolytic activity.

- 5 23. A method of treating a patient in need of such treatment for a condition which is mediated by protease D-G, comprising administration of a protease D-G modulating compound active in the method of Claim 16.